



## Louisiana Envirothon Training

# Section 1: Mammal Skulls



# Identifying Mammal Skulls

The first step to identifying an animal skull is to distinguish certain features generally referred to as Best Recognition Factors (BRF).

BRF's of mammal skulls can help you identify which class (herbivore, carnivore, omnivore) and species of animal the skull belongs to...



**A good place to start is with the  
teeth of the skull.....**

**The types of teeth you see will  
help determine if the animal is  
a carnivore, an omnivore or an  
herbivore.**




## Mammal Teeth (4 main types)

**Incisors:** located at the front of the mouth with a sharp, flat biting surface; used for cutting, shearing, and pulling food

**Canines:** long, curved, and pointed; used for piercing and tearing food

**Premolars:** flat biting surface; may have a cusp along the edge; used for tearing and crushing food

**Molars:** flat biting surface; may have cusps at the edges; used for chewing, crushing, and grinding food



[www.nps.com](http://www.nps.com)



# Mammal Teeth

## Carnivores: primarily meat eaters

- Teeth are designed for cutting, tearing, and piercing
- Small incisors
- Large, sharp canines
- Premolars and molars are used for shearing and will have sharp cusps

## Herbivores: primarily plant eaters


- Teeth are designed for breaking down tough plant foods
- Wide incisors are used for stripping or cropping vegetative matter
- Canines are small or absent
- Premolars and molars of grazers are very flat
- Premolars and molars of browsing animals have low, sharp cusps

## Omnivores: eat both meat and plants

- Combination of herbivore and carnivore teeth
- Long, sharp canines, but not as long as carnivores
- Premolars have sharp cusps
- Molars are usually squared off and bumpy



**Carnivores** tend to have long canines which are used to rip and tear meat, sometimes in a scissors like action. In addition, carnivores have sharp molars toward the back of the mouth, used to further rip and shred meat. Carnivores tend to have binocular vision, where their eyes are at the front of the head, which results in a smaller field of view, but allows for depth perception, needed to catch prey. Examples of carnivores include otters and bobcats.



River Otter Skull




Text Taken From: <https://www.chesco.org/DocumentCenter/View/47500/2019-What-Can-I-Learn-From-a-Skull?bidId=>

Photo From Skulls Unlimited

**Herbivores** tend to have well-developed flat premolars and molars, often with sharp ridges on the tops. Generally herbivores do not have canine teeth, and their incisors are usually large and used to snip off foliage from branches. Because herbivores are often prey for other animals, they generally have their eyes on the side of their head, which functions to give them a wider field of view, so that they can detect their prey earlier, and have a chance to flee. Examples of herbivores include rabbits, beaver, nutria, muskrat, and white-tailed deer.

White-tailed deer Skull




Text Taken From: <https://www.chesco.org/DocumentCenter/View/47500/2019-What-Can-I-Learn-From-a-Skull?bidId=>

Photo From Acorn Naturalists



**Omnivores** usually have a variety of all kinds of teeth. Generally omnivores have eyes on the front of their heads like carnivores, in order to best catch their prey. Examples of herbivores include black bear, raccoons, coyote, fox and opossums.

Virginia Opossum Skull



Text Taken From: <https://www.chesco.org/DocumentCenter/View/47500/2019-What-Can-I-Learn-From-a-Skull?bidId=>

Photo From Skulls Unlimited

# Mammal Nose and Eye Features Can Also Help To Identify Skull Species

## The Rostrum: nasal area

- Size is **related to importance of smell**
- **Herbivores generally have long noses**
- Noses of omnivores and carnivores vary

Eyes in the front, the animal hunts. Eyes on the side, the animal hides.

## **Orbit:** part of skull surrounding the eyeball

- **Carnivores have large forward facing orbits allowing for binocular vision and depth perception**
- **Herbivores have eyes on the side allowing sight in all directions (almost 180 degrees of vision per side)**
- **Grazers (eat low lying vegetation) tend to have eyes high on the head to see above grass (ex. Cows)**
- **Browsers (eat high growing plants) have eyes far back on the skull to avoid twigs and preserve vision when their noses are in brush (ex. Deer)**
- **Omnivores tend to have orbits facing partially forward giving maintaining a wide field of view and decent depth perception**
- **Animals that spend time in the water tend to have orbits high on the skull to the eyes above water**
- **Nocturnal animals tend to have large eyes and orbits for increased night vision**



# The Skulls Of Louisiana Mammals You Are Learning To Recognize Are:

- Louisiana Black Bear
- Feral Hog
- Fox-Gray and Red
- Coyote
- Bobcat
- (Virginia) Opossum
- (American) Beaver
- Nutria
- Muskrat
- (American) River Otter
- (American) Mink
- Raccoon
- (Nine banded) Armadillo





# Louisiana Black Bear Skull



Photo from Skulls Unlimited



# Feral Hog Skull



Photo from Skulls Unlimited





# Red Fox and Gray Fox

Grey Fox

Red Fox

Ridges form a 'U'

Ridges form a 'V'



Red Fox



Gray Fox



Coyote: shown for reference



# Coyote Skull



Photo from Skulls Unlimited







# Bobcat Skull



Photo from Skulls Unlimited







# Virginia Opossum Skull



Photo from Skulls Unlimited





# American Beaver Skull



Photo from Skulls Unlimited





# Nutria Skull



Photo from Skulls Unlimited





# Muskrat Skull



Photo from Skulls Unlimited





# American River Otter Skull



Photo from Skulls Unlimited





# American Mink Skull



Photo from Skulls Unlimited





# Raccoon Skull



Photo from Skulls Unlimited







# Nine Banded Armadillo Skull



Photo from Skulls Unlimited






# Louisiana Furbearer Pelts







# Bobcat




Bobcats live in a vast variety of habitat types, including heavily forested areas, swamps, bottomland hardwoods, and even deserts.



**Males weigh 20 to 22 pounds; females weigh 18 to 19.**



# Coyote




**Coyotes live in a vast variety of habitat types, including forested areas, farmlands, prairies swamps, mountains, and deserts.**



**Males average 30 pounds; females 25. Coat color varies, but generally mottled gray with a lighter belly.**



# Red Fox



Red foxes live in mixed wooded areas with ungrazed pastures, rice fields, and cane fields.




**Adult weight is 8-14 pounds and the total length is 3-4 feet (12-17 inches of which is tail). They have yellow eyes with elliptical pupils.**








# Mink




Mink live along ponds, canals, rivers, streams, lakes and marshes.



**Males can be 20 to 30 inches and weigh over three pounds; females are 1 ½ to 2 pounds and 16 to 21 inches.**



# North American River Otter



Otters live near streams, lakes and marshes. They prefer isolated habitats.




**Males can weigh up to 25 pounds; females generally weigh under 20.**





# North American Raccoon




Raccoons hunt in and around water, but they can roam far from water. They can often be found living in urban habitats.



**The hind legs are longer than the front legs so that they have a hunched posture.**



# Striped Skunk




Striped skunks prefer a mixture of woodlands and farmlands.



**Skunks can spray their musk repeatedly when they feel threatened.**



# North American Beaver



Beavers live near wooded rivers, streams, lakes, swamps, and backwaters.




**Beavers have a humped back with a wide flat tail and webbed feet. The average weight is 33 pounds.**







# Nutria



Nutrias live in swamps, marshes, rivers, lakes, streams, back waters.




**Nutria are a non-native furbearer. They were imported from South America. They are considered an invasive species.**

**Nutria have mammary glands on the sides so that infants can nurse while the mother swims.**



# Virginia Opossum



**Opossums adapt to a variety of habitats: farmlands, prairies, swamps, and forests**



**Opossums have more teeth than any other mammals. They are the only marsupial in North America.**





# Size and Shape comparison of water mammals



**Mink**



**Muskrat**



**Nutria**



**Beaver**



**Otter**



Otter

Mink

Beaver

Nutria

Muskrat



# Section 2: General Biology






# Definitions

- Environment – a physical location in time and space, small or large, stable or temporary including both the physical conditions (Abiotic) and organisms (Biotic)
- Abiotic – non-living components of an ecosystem (weather, soil, water)
- Biotic – living components of an ecosystem (animals, plants, fungi)
- Ecosystem – Biotic and abiotic components of an environment functioning together as a system



# Ecosystem of a forest



# Definitions

- **Habitat**: the place where a particular plant or animal is naturally found.
- The animals physical needs determine where a species will exist. Habitats provide energy (in the form of food and water) and shelter
  - Examples of habitat factors can include: Vegetation type, weather, amount/type of water present, food & prey/predator availability. Successful reproduction
- Animals are designed for or adapted to the habitats where they live. That is, they have developed bodies and behaviors to help them survive in their environment. **Adaptations** help animals get food, protect themselves, and reproduce.



Adaptations genetically determined characteristics of an animal that help it to survive in its environment. These characteristics fall into three main categories: body parts, body coverings, and behaviors. Any or all of these types of adaptations play a critical role in the survival of an animal.

### Physical adaptations

- **Type of body covering:** fur, feathers, scales
- **Color:** patterns, match surroundings
- **Body part:** beak, antlers, ears, claws
- **Defenses:** venom, spray, quills



# Examples of animal adaptations include:

- the long necks of giraffes for feeding in the tops of trees
- the streamlined bodies of aquatic fish and mammals
- the light bones of flying birds and mammals
- and the long dagger-like canine teeth of carnivores



## Some Bat Adaptations



- Light bones for flying
- Big ears for Eco locating
- Specialized mouth for capturing prey






# Food Chains and Food Webs

A food chain is a possible path that energy and nutrients may take as they move through the ecosystem. All food chains start with a producer (a plant), which is eventually eaten by a consumer.



# Food Chains and Food Webs

A food web consists of **all the food chains** in a **single ecosystem**. All of the interconnected and overlapping food chains in an ecosystem make up a food web.



# Section 3: Biodiversity and Endangered Species





**Biodiversity is the variety of different types of life found on earth. It is a measure of the variety of organisms present in different ecosystems.**



**The more plant, insect, and animal species there are in one area the greater the biodiversity and generally the healthier the ecosystem.**



**All living creatures need other creatures and plants in one way or another even if the connection is not so clear.**

**Therefore the removal of a single species can conceivably set off a chain reaction affecting many others.**

**Due to this fact, a law was created to help prevent the loss of any species of animal or the habitat they live in the United States.**







# The Endangered Species Act of 1973



The Endangered Species Act (ESA) provides a program for the conservation of threatened and endangered plants and animals and the habitats in which they are found. The United States Fish and Wildlife (USFWS) enforces regulations pertaining to this act.

Red cockaded woodpecker and long leaf pine habitat  
USFWS Photos



# Key Definitions

**Endangered:** Any species in danger of extinction throughout all or a significant portion of its range.

**Threatened:** Any species likely to become endangered within the foreseeable future throughout all or a significant portion of its range.

**Conserve:** Use of all methods and procedures necessary to bring any endangered or threatened species to the point at which the measures provided under the Act are no longer necessary.

**Keystone species:** A species on which other species in an ecosystem largely depend, such that if it were removed the ecosystem would change drastically.

Examples include: sharks, grizzly bears, wolves, sea otters, prairie dogs and snowshoe hares



Western Indian Manatee

USFWS photo



# What Causes Animals to Become Endangered?

- **Loss of Habitat** – The most significant causes of endangered animals is habitat loss
- **Invasive Species** – Introduced species generally outcompete native species for resources
- **Overexploitation of Resources** - Human actions that directly cause a decline in a species population (ex: overfishing, historical collection of species)
- **Pathogens and Disease** - native populations have little resistance to the invading pathogen introduced into their environment by invasive species or humans.
- **Environmental Pollution** – addition of pollutions and toxins into a habitat that negatively affect a species (ex: pesticides including DDT)



Photo of habitat loss  
from Wikipedia

# Section 3: Endangered Species in Louisiana

## Whooping Crane



Whooping Cranes are one of the worlds most endangered crane species with an estimated 700 individuals in the world.

They only occur naturally in North America. Louisiana began a project in 2011 attempting to reestablish a population of whooping cranes in Louisiana. As of October 2021, the state has 71 cranes.

They historically utilized the Cajun Prairie grasslands, however, now they most commonly occur in rice and crawfish fields.

One of the biggest issues LA's current Whooping Crane population faces is awareness and proper identification. Residents struggle to identify the crane when compared to other egret species we have in the state.





Photo by Charles Martin



Whooping Crane

## Several Species of Birds Resemble Whooping Cranes

The *Snow Goose* is the **ONLY** species that can be shot during hunting season.

**KNOW YOUR TARGET BEFORE YOU SHOOT!**

Photo by Simon Pierre Busetta, Wikimedia Commons



Snow Goose  
*\*Only LEGAL game species\**

Photo by Eric Uffman



White Ibis

Photo by Steve Uffman



Wood Stork

Photo by Steve Uffman



Great Egret

Photo courtesy of USFWS



Sandhill Crane

Photo by Steve Uffman



White Pelican



# Size comparison



Great Egret



Snowy Egret



This is a Whooping Crane





# Whooping Crane is North America's Tallest Bird


Height: 5 feet  
Weight: 5-7 kg (11-17 lbs)  
Wingspan: ~7-8 ft



# A few of Louisiana's other Endangered Species include:

- Red Cockaded Woodpecker
- Western Manatee
- All sea turtle species
- Louisiana Pine Snake
- Abbeville Red Iris
- Gulf Sturgeon

The complete list can be found at [here](#) and [here](#)





# The Success Stories

Louisiana has been successful in recovering several species that were once considered threatened or endangered. These animals are more numerous now.

Examples of Louisiana success stories include: the Brown Pelican, the Louisiana Black Bear, the Bald Eagle and the American Alligator.



Photo by Heather Williams





# Section 4: Invasive Species in Louisiana



# Key Definitions

Invasive species is an introduced organism that becomes overpopulated and negatively alters its new environment.

Examples of terrestrial invasive species in Louisiana are:

Feral hog

Imported Red fire ant

Asian tiger mosquito

Nutria



# Why Are Invasive Species Bad?

**Invasive species are harmful in Louisiana because they may:**

- Spread rapidly due to a lack of natural predators
- Feed on native species and reduce native population sizes
- Outcompete native species for food and habitat resources
- Introduce new parasites and pathogens

**A species is considered invasive when:**

1. It is introduced into a new area, outside of its native range
2. Its presence disrupts native food webs



Louisiana's sub-tropical climate provides suitable habitat for invasive species from all over the world. Stopping the spread of aquatic invasive species is extremely difficult, especially in Louisiana with our interconnected waterways. LDWF needs your help to locate invasive species and to prevent new invasive species from becoming established.

*A waterbody plagued by  
Silver Carp.*

Nerissa Michaels/AP/Nicola's River Biological Station via  
HoumaToday.com

## How Can You Help?

Information about the location of various invasive species is an important tool in helping biologists combat spreading. Fortunately, you can help by taking the following actions:

**If you see/catch any of these species:**

- Record date and location (GPS coordinates preferred)
- If caught, place organism in a labeled bag and freeze
- Contact LDWF at 225-765-0765

**Other ways to help:**

- Never release aquarium pets purchased from pet stores into the wild. Contact LDWF for proper disposal of unwanted pets.
- Never empty live fish, bait, or water from a bait well from one body of water into another.



**Invest in the Future...Geaux Fish Louisiana!**

This public document was published at a total cost of \$\_\_\_\_\_ copies of this public document were published and partially paid for by an Aquatic Invasive Species Grant from the U.S. Fish and Wildlife Service. In this first printing at a cost of \$\_\_\_\_\_. This document was published by the Louisiana Department of Wildlife and Fisheries, 3000 Quail Drive, Baton Rouge, LA 70808, to inform Louisiana residents and non-residents about the impacts of aquatic invasive species. This material was printed in accordance with standards for printing by state agencies.



# BEWARE

## Saltwater

### Asian Tiger Prawn

(*Penaeus monodon*)

- Dark color with white and yellow stripes
- Occasionally have red line extending length of body
- Larger than and competes with native shrimp



### Lionfish

(*Pterois spp.*)

- Red, white, and brown stripes with elongated dorsal and pectoral fins
- Found on reef structures and hard bottoms
- Venomous spines
- Aggressive predator that competes with and feeds on native reef organisms



## Freshwater

### Northern Snakehead

(*Channa argus*)



- Brown with dark blotches, long dorsal and anal fins, flattened head
- Aggressive predator
- Often confused with the native bowfin

### Rio Grande Cichlid

(*Herichthys cyanoguttatum*)



- Dark color, white and blue spots
- Large forehead
- Territorial
- Competes with and often confused with native panfish

### Apple Snail

(*Pomacea spp.*)



- Brown, black, or yellow banded shell
- Large, up to 10 cm
- Attach pink or orange egg clusters to vegetation or structures above water line
- Feeds on aquatic vegetation
- May carry parasites that infect humans

### Silver Carp

(*Hypophthalmichthys molitrix*)



- Leaps out of water and injure boaters
- Competes with native filter feeding fish & shellfish

### Bighead Carp

(*Hypophthalmichthys nobilis*)



- Competes with native filter feeding fish & shellfish

### Common Carp

(*Cyprinus carpio*)



- Disturbs bottom sediments and uproots native aquatic vegetation when feeding

### Black Carp

(*Mylopharyngodon piceus*)



- Threatens populations of native snails and mussels

### Grass Carp

(*Ctenopharyngodon idella*)



- Disturbs bottom sediments and uproots native aquatic vegetation when feeding

# Aquatic Invasive Species

Controlling and Identifying  
Aquatic Invasive Species in Louisiana

[www.wlf.la.gov](http://www.wlf.la.gov)





# BEWARE

Water Hyacinth



*Eichhornia crassipes*

**Origin: Amazon Basin, South America**

- Free floating
- Leaves shiny green, leathery, and oval with gently incurved sides that stand erect
- Leaf veins are dense and numerous
- Spongy thick stalks
- Purple flowers with six petals
- Roots are dark and feathery

Giant Salvinia



*Salvinia molesta*

**Origin: Brazil**

- Free floating
- Small, oblong, spongy, green leaves in whorls of three; two floating and one submerged
- Margins of mature plants curl inward
- Leaf surface has rows of hairs that, when magnified, are eggbeater shaped. Hairs give the leaves a velvety appearance and repel water
- No flowers

Water Lettuce



*Pistia stratiotes*

**Origin: Amazon Basin, South America**

- Free floating
- Resembles a floating open head of lettuce
- Leaves are thick, hairy, ridged, and pale green
- Leaf margins are wavy (with top margins scalloped)
- No leaf stems
- Flowers are inconspicuous
- Fruit is a green berry
- Numerous feathery roots hang submerged beneath leaves

Hydrilla



*Hydrilla verticillata*

**Origin: China/Asia**

- Submerged perennial
- Stems are slender and branched, covered with small pointed, often serrated, leaves arranged in whorls of four to eight
- Leaf midribs are often reddish with one or more sharp spines
- Branching stems reach the surface and form dense mats
- Small white flowers

Common Salvinia



*Salvinia minima*

**Origin: South and Central America**

- Free floating fern
- Similar in appearance to giant salvinia, except has stiff leaf hairs with branches free at the tips
- Root-like structures hang below the surface
- No flowers

Alligator Weed



*Alternanthera philoxeroides*

**Origin: South America**

- Emerged perennial
- Long, branched, hollow stems
- Leaves are simple, elliptic, and opposite with smooth margins
- Solitary white flowers that grows on stalks during warm months
- Fibrous roots

## Aquatic Invasive Plants

Identifying and Preventing  
Aquatic Invasive Plants in Louisiana

[www.wlf.la.gov](http://www.wlf.la.gov)





# Aquatic Invasive Plants Can Cause Serious Environmental and Economical Harm

## Impacts

### Reduce or eliminate native fish, bird, and plant populations

- Out-compete native plants
- Alter fish habitat
- Reduce waterfowl habitat and food

### Clog waterways

- Limit boating and fishing access

### Alter water quality

- Cause changes in pH
- Reduce dissolved oxygen
- Increase temperature

A Louisiana waterbody plagued by giant salvinia.



## Control Efforts

There are various control methods that have been used to combat the spread of invasive plant species. Each one has pros and cons in its ability to treat infested waterbodies.

	Pros	Cons
<b>CHEMICAL</b>		
Herbicides	✓ Quick treatment, effective, can improve fish habitat	✗ Costly; weather dependent
<b>BIOLOGICAL</b>		
Alligator Weed Flea Beetle	✓ Can thin and limit alligator weed growth to allow navigation in previously inaccessible areas	✗ Localized effectiveness only during parts of growing season
Salvinia Weevil	✓ Species-specific insects that feed on the buds of common and giant salvinia and whose larvae tunnel into the rhizomes of the plant causing it to die and sink	✗ This is a long-term control measure that can take several years; very intolerant of cold temperatures
Triploid Grass Carp	✓ Sterile, non-reproducing grass carp which consume submerged aquatic vegetation	✗ Not specific in the vegetation it consumes; older, bigger fish are not as effective
<b>MECHANICAL</b>		
Drawdowns	✓ Cheap, effective, and can improve fish habitat	✗ Require dry conditions and long time frame; restricts boating access
Physical Removal	✓ Can target specific areas; available on windy days or when other options are not available	✗ Extremely labor intensive

## How Can You Help?

Preventing new introductions is the best and most cost-effective way to control the impacts of invasive species. Fortunately, there are a few simple actions you can take to prevent their spread:

**INSPECT** your boat, trailer, and equipment. Remove any visible plants, animals, or mud before leaving a waterbody.

**REMOVE** unwanted bait from boat, live well, or bucket. **DISCARD IN TRASH, NOT WATER!**

**DRAIN** water from your boat, motor, live well, and bilge before leaving a waterbody.


**RINSE** and **DRY** your boat, trailer, and fishing equipment to remove or kill species that were not visible when you left a waterbody. Before going to another water body, rinse boat and trailer with hot (104 degrees) water or allow boat and trailer to dry for at least five days.

Learn to identify aquatic nuisance species and report any new infestations to the Louisiana Department of Wildlife and Fisheries at (225) 765-2328.

**Invest in the Future...Geaux Fish Louisiana!**

This public document was published at a total cost of \$\_\_\_\_\_ copies of this public document were published and paid for by the Aquatic Plant Fund in this first printing at a cost of \$\_\_\_\_\_. This document was published by the Louisiana Department of Wildlife and Fisheries, 2000 Quail Drive, Baton Rouge, LA 70806, to inform Louisiana residents and non-residents about the Sport Fish Restoration Program. This material was printed in accordance with standards for printing by state agencies.

# Section 6: Managing For Wildlife



A wildlife management plan is a description of the short-term objectives and long-term goals that will be met by manipulation of habitat, wildlife populations, and people. Management plans explains how these objectives and goals can be reached for a specific location. All species need food, cover (or shelter), space, and water. A wildlife management plan must address all four of these needs.

Wildlife management takes into consideration the ecological principles such as carrying capacity of the habitat.

The carrying capacity of an environment is the maximum population size of a biological species that can be sustained by that specific environment, given the food, habitat, water, and other resources available.





## Common wildlife management practices:

- Hunting
- Trapping
- Selective cutting of trees opens the canopy of the forest, allowing the understory to grow
- Fires and selective burning
- Food plots



# Common wildlife management practices continued:

- Conservation easements (agreement between a landowner and a land trust or government agency that permanently limits uses of the land in order to protect its conservation values)
- Wildlife refuges and wildlife management areas
- Creating snags
- Controlling invasive plants

Washing feet and fish nets can help prevent the spread of invasive species such as Salvinia and Water Hyacinth



Which of these and other management practices are applied will depend upon what species or habitat type you are managing for.



If you are a property owner and you want to manage for a certain species or habitat type, you would contact different agencies for assistance. Here in Louisiana, the Department of Wildlife and Fisheries (LDWF) (state agency) or the USFWS (federal agency) can help.

### LDWF

- Helps to protect all wildlife and all aquatic life within the state of Louisiana. LDWF specializes in assisting with game and nongame species management. So for example, if you wanted assistance in modifying habitat for deer, quail, or songbirds, you could ask for assistance from LDWF.

### USFWS

- Helps to protect and recover threatened and endangered species and monitor and manage migratory birds. So for example, if you wanted assistance in modifying habitat for Gopher Tortoises or Red-cockaded Woodpeckers, you could ask for assistance from the USFWS.

